BUTTOCK LIFT (BBL)

What To Expect

After Surgery

Returning To Daily Activities

These instructions are provided to assist in the healing process and ensure best results. If you choose not to follow these instructions, it may increase your risk of possible complications and compromise results of the procedure.

3

ficlinica

What To Expect After Surgery

HOSPITAL RECOVERY & DISCHARGE

- ⁷After a buttock lift surgery it is normal to stay in the clinic for 1 night. You will be offered a menu to select meals and beverages for the duration of your stay.
- [/]After your surgery you will wake up in your room dressed in the compression garment. Likely you will have drains. Typically, drains are removed on the day of discharge.
- ⁷The medical staff will regularly monitor you during your stay in the clinic. We encourage you to maintain open communication with them to help make your stay as comfortable as possible.
- [/]Upon discharge you will be provided with a "goodie bag" with all essential medications, antibiotics and dressings for your recovery at home as well as written post-operative instructions.

PAIN LEVEL

Expect to have pain and numbress in areas. Maximum discomfort will occur during the first few days. Pain should be controlled by painkillers. For dosage please refer to the instructions leaflet in the "goodie bag".

COMPRESSION GARMENT

⁷You are required to wear the supplied garment for 23hrs a day for 6-8 weeks. If you choose to buy a spare compression garment, ensure it is the exact same one. Using a different garment than the one provided by the clinic may affect your surgery results and your eligibility for corrective surgery.

SUTURES

Your incision will have internal sutures that will dissolve in 6 months.

DRESSING CHANGE

⁷Dressing should be changed every 2-3 days or more often if there is a lot of discharge from the wound. Nurses will do your first dressing change in the clinic. Pay attention to dressing change steps, make notes in case you will need to instruct your partner back home.

Buttock Lift (BBL)

SHOWER

⁷Use sensitive cleansing wipes to save you from extra trips to the bathroom. You can take a shower 3-4 days after surgery. Let soapy water run over incisions, but do not scrub them.

COPING EMOTIONALLY

'Surgery places stress on the body and mind. Emotional stress is a real thing in the post-operative period. You may second guess your decision to undergo surgery, you may have feelings ranging from anger to guilt, you may even find yourself in tears just wanting to feel like your normal self again. This is natural phase of healing process and will soon becomes a distant memory.

POST OP APPOINTMENTS

Post-op appointments can be booked at the front desk or by contacting the clinic. At the Kaunas clinic, post-op check-ups are free within the first year after surgery. In London and Dublin, appointments are available according to our visit schedule, with a non-refundable fee. Due to limited availability, advance registration and payment are required.

SCAR CARE

Scar care is about helping your skin heal well and minimizing the scar's appearance over time. It involves keeping the scar clean, moist, and protected. Use silicone gel or sheets to help flatten and soften scars, massage the area gently to improve circulation, and always protect it from the sun with sunscreen or clothing to prevent darkening. Consistency is key for the best results.

Returning To Daily Activities

HEALING & SWELLING

- You will experience swelling and bruising. The majority of this will subside in 6-8 weeks. Your recovery is subject to your response to the healing process and is different for everyone. Skin firmness, hypersensitivity, or lack of sensitivity is is normal and will gradually resolve over time.
- ⁷Sleep on your stomach or side to avoid pressure on your buttocks. Use pillows for support and alignment. Avoid sitting for long periods.
- ⁷After surgery, you'll usually need a series of manual lymphatic drainage massages. The number needed varies, but it's best to start with frequent sessions and reduce them as swelling and

Buttock Lift (BBL)

healing improve. Be sure to choose an experienced physiotherapist.

- ⁷ It is quite common for your sides to heal differently. It is normal if one may swell more than the other or heal slower than another. Fight the temptation to compare yourself to people across the internet. Your journey is unique and valid. Everyone's situation is different. Be patient as you heal your real results won't reveal themselves for at least 4–6 months, so don't panic!
- ⁷Do not touch the incision sites. If this is unavoidable, make sure to disinfect hands and incision area first.
- ⁷Surgical procedures can often dry the skin, and compression garments may add to this by rubbing against the skin. To help maintain healthy skin around the incision area, we recommend moisturizing thoroughly. Keeping the area well-moisturized can aid in your comfort and support the healing process.

WARNINIG SIGNS

- [/]A fever of 38°C or higher.
- Shaking or chills.
- [/]Heat in the area and bright red skin that is hot to the touch.
- [/]Nausea or vomiting.
- [/]Increased pain, swelling and redness that's getting worse.
- [/]Continuous fluid, pus and smell from your stitches.
- [/]Separation of your stitches.
- [/]Sudden shortness of breath or trouble breathing.
- ⁷Any unexplained or unexpected symptoms and any pain that cannot be controlled by your pain medication.

HYDRATION & NUTRITION

- Rehydrate your body often. Surgery of any kind can reduce fluids in your body. Drinking water frequently will help replenish these lost body fluids. Also, eat lightly for the first few days and only food that is easy to swallow and digest.
- ⁷Do not drink any alcohol if you are taking any medication.

EXERCISE & ACTIVITY

⁷After surgery you will be mobile. Start gentle walking as soon as possible to help reduce swelling and lower the risk of blood clots.

Buttock Lift (BBL)

- Please refrain from housework and lifting and stretching for minimum of 4 weeks.
- Strenuous exercise and activities are restricted for 8-12 weeks. It may cause increased swelling, bleeding, or disruption of the healing process.

SWIMMING & SUN EXPOSURE

- No bath, pool, jacuzzi, ocean water for 8-12 weeks post-surgery or until your wounds are fully healed.
- Your scars may remain red and raised for several months. Most patients notice them gradually fading and softening over the course of 12–18 months, but this process can vary. Be patient—scar improvement takes time. Avoid exposing scars to the sun and sunbedding. If sun is unavoidable always apply strong sunscreen.
- We advise waiting for at least 6 weeks before a spray tan appointment or until the incision heals completely.

DRIVING

⁷You are not allowed to drive for the first few weeks after your operation for safety reasons. To be fit to drive, you must no longer be taking any sedative medications and must be able to operate your car safely and comfortably. For most people to be able to react and move without pain would be around 3-6 weeks. Swelling and pain may increase after driving.

MAINTAINING RESULTS

⁷Every patient is unique and surgery results can vary from patient to patient. Your body will change with age. The appearance of your buttocks will change too. To maximise the longevity of your surgery results, it is important to maintain a healthy lifestyle. Please keep in mind that your surgery results may be impacted by pregnancy, fluctuations in your weight, lifestyle and overall health.

RETURNING TO WORK

⁷ Each patient is different, so your time off work will vary depending on your recovery speed and the type of work you do. Allow yourself at least 2 weeks to recuperate before returning to non-strenuous work. Remember, it is up to you to decide when you are ready to return. If your job is physically demanding, you may need to take more time off.

Tell Your Story

Join our Facebook group BEFORE & AFTER Fi Clinica Europe


Your feedback inspiration for others!












booking@ficlinica.com



@ Ficlinica